


# IOQM 2022 Part B

## Official Solutions

**Problem 1.** Let  $D$  be an interior point on the side  $BC$  of an acute-angled triangle  $ABC$ . Let the circumcircle of triangle  $ADB$  intersect  $AC$  again at  $E(\neq A)$  and the circumcircle of triangle  $ADC$  intersect  $AB$  again at  $F(\neq A)$ . Let  $AD$ ,  $BE$  and  $CF$  intersect the circumcircle of triangle  $ABC$  again at  $D_1(\neq A)$ ,  $E_1(\neq B)$  and  $F_1(\neq C)$ , respectively. Let  $I$  and  $I_1$  be the incentres of triangles  $DEF$  and  $D_1E_1F_1$ , respectively. Prove that  $E, F, I, I_1$  are concyclic.



**Solution.** Note that

$$\angle C F_1 D_1 = \angle C A D_1 = \angle E A D = \angle E B D = \angle E_1 B C = \angle E_1 F_1 C,$$

so  $F_1 C$  is the bisector of  $\angle D_1 E_1 F_1$ . Similarly,  $E_1 B$  is the bisector of  $\angle D_1 E_1 F_1$ , implying  $I_1 = BE_1 \cap CF_1$ . Now,

$$\begin{aligned} \angle EDF &= \angle EDA + \angle FDA = \angle EBA + \angle FCA \\ &= \angle E_1 BA + \angle F_1 CA = \angle E_1 D_1 A + \angle F_1 D_1 A = \angle E_1 D_1 F_1. \end{aligned}$$

Therefore

$$\angle EIF = 90^\circ + \frac{1}{2} \angle EDF = 90^\circ + \frac{1}{2} \angle E_1 D_1 F_1 = \angle E_1 I_1 F_1 = \angle EI_1 F,$$

which proves the required concyclicity.  $\square$

**Problem 2.** Find all natural numbers  $n$  for which there exists a permutation  $\sigma$  of  $1, 2, \dots, n$  such that

$$\sum_{i=1}^n \sigma(i)(-2)^{i-1} = 0.$$

*Note:* A permutation of  $1, 2, \dots, n$  is a bijective function from  $\{1, 2, \dots, n\}$  to itself.

**Solution.** Suppose that  $n \equiv 1 \pmod{3}$  and  $\sigma$  a permutation of  $1, 2, \dots, n$ . Then

$$\sum_{i=1}^n \sigma(i)(-2)^{i-1} \equiv \sum_{i=1}^n \sigma(i) = \frac{n(n+1)}{2} \pmod{3},$$

and hence the left-hand side is non-zero.

We now show by induction that if  $n \equiv 0$  or  $2 \pmod{3}$  then there exists a permutation of  $1, 2, \dots, n$  satisfying the given condition.

If  $n = 2$  then the permutation given by  $\sigma(1) = 2, \sigma(2) = 1$  satisfies the given condition. Similarly, if  $n = 3$  then the permutation  $\sigma(1) = 2, \sigma(2) = 3, \sigma(3) = 1$  satisfies the given condition.

Suppose that for  $n = m$  there exists a permutation  $\sigma$  satisfying the given condition. We consider the permutation  $\tau$  of  $1, 2, \dots, m+3$  given by  $\tau(1) = 2, \tau(2) = 3, \tau(m+3) = 1$  and  $\tau(i) = \sigma(i-2) + 3$  for  $i = 3, 4, \dots, m+2$ . Then


$$\begin{aligned} \sum_{i=1}^{m+3} \tau(i)(-2)^{i-1} &= 2 - 6 + (-2)^{m+2} + \sum_{i=3}^{m+2} 3 \cdot (-2)^{i-1} \\ &= 2 - 6 + (-2)^{m+2} - 4 \cdot ((-2)^m - 1) = 0. \end{aligned}$$

Thus, by induction it follows that for every  $n \equiv 0$  or  $2 \pmod{3}$  there exists a permutation satisfying the given condition.  $\square$

**Problem 3.** For a positive integer  $N$ , let  $T(N)$  denote the number of arrangements of the integers  $1, 2, \dots, N$  into a sequence  $a_1, a_2, \dots, a_N$  such that  $a_i > a_{2i}$  for all  $i$ ,  $1 \leq i < 2i \leq N$  and  $a_i > a_{2i+1}$ , for all  $i$ ,  $1 \leq i < 2i+1 \leq N$ . For example,  $T(3)$  is 2, since the possible arrangements are 321 and 312.

- (a) Find  $T(7)$ .
- (b) If  $K$  is the largest non-negative integer so that  $2^K$  divides  $T(2^n - 1)$ , show that  $K = 2^n - n - 1$ .
- (c) Find the largest non-negative integer  $K$  so that  $2^K$  divides  $T(2^n + 1)$ .

**Solution.** (a) Given an arrangement  $a_1, a_2, \dots, a_7$ , satisfying the given conditions, we can build a binary tree with nodes as in the Figure below. At each node, the root node



is greater than the child nodes. Conversely, any such tree gives a valid arrangement. Observing that the root of the tree must contain the maximum of the numbers, we can choose 3 out of the other 6 numbers in  $\binom{6}{3}$  ways and build the left tree and the right tree, each in 2 ways and hence the number of such trees is  $2 \cdot 2 \cdot \binom{6}{3} = 80$ .

(b) Observe that  $T(N)$  is also the number of ways of arranging any  $N$  distinct numbers into a sequence  $a_1, a_2, \dots, a_N$  satisfying the given conditions. Also, the given conditions imply that  $a_1 = \text{maximum of the numbers}$ . Now, leaving out the maximum, the rest of the  $2^n - 2$  numbers can be split into two groups of  $2^{n-1} - 1$  numbers each and these can be individually put into a sequences  $b_1, b_2, \dots, b_{2^{n-1}-1}$  and  $c_1, c_2, \dots, c_{2^{n-1}-1}$  satisfying the

conditions in  $T(n - 1)$  ways each. Now, the required arrangement of the original given sequence can be obtained as follows:

$$a_1, b_1, c_1, b_2, b_3, c_2, c_3, b_4, b_5, b_6, b_7, c_4, c_5, c_6, c_7, \dots$$

This gives

$$T(2^n - 1) = T(2^{n-1} - 1)^2 \binom{2^n - 2}{2^{n-1} - 1} \quad (1)$$

We find the highest power of 2 that divides  $\binom{2^n - 2}{2^{n-1} - 1}$ :

We have

$$\begin{aligned} 2^{n-2} \binom{2^n}{2^{n-1}} &= 2^{n-2} \cdot \frac{2^n!}{2^{n-1}! 2^{n-1}!} \\ &= 2^{n-2} \cdot \frac{2^n(2^n - 1)(2^n - 2)!}{2^{n-1}(2^{n-1} - 1)! 2^{n-1}(2^{n-1} - 1)!} \\ &= (2^n - 1) \binom{2^n - 2}{2^{n-1} - 1} \end{aligned}$$

Now, the highest power of 2 that divides  $\binom{2^n}{2^{n-1}}$  is

$$(2^{n-1} + 2^{n-2} + \dots + 1) - 2(2^{n-2} + 2^{n-3} + \dots + 1) = 1$$

Hence the highest power of 2 that divides  $\binom{2^n - 2}{2^{n-1} - 1}$  is  $n - 1$ .

From the recurrence (1), if  $t_n$  is the highest power of 2 dividing  $T(2^n - 1)$ , then  $t_n = 2t_{n-1} + n - 1$ . From the initial conditions,  $t_1 = 0, t_2 = 1, t_3 = 4$ , we obtain, by an easy induction, that  $t_n = 2^n - n - 1$ .

(c) Suppose that  $N = 2^n + 1$ . It is easy to see that

$$T(2^n + 1) = T(2^{n-1} - 1)T(2^{n-1} + 1) \binom{2^n}{2^{n-1} + 1}$$

The highest power of 2 dividing  $\binom{2^n}{2^{n-1} + 1}$  is  $n$ :

$$(2^{n-1} + 1) \binom{2^n}{2^{n-1} + 1} = \binom{2^n}{2^{n-1}} \cdot 2^{n-1}$$

Since the highest power of 2 dividing  $\binom{2^n}{2^{n-1}}$  is 1, it follows that the highest power of 2 dividing  $\binom{2^n}{2^{n-1} + 1}$  is  $n$ . Thus, if  $s_n$  denotes the highest power of 2 dividing  $T(2^n + 1)$ , then

$$s_n = s_{n-1} + 2^{n-1} - (n - 1) - 1 + n = s_{n-1} + 2^{n-1}$$

Hence  $s_n - s_1 = 2^n - 2$  and since  $s_1 = 1$  (since  $T(3) = 2$ ), it follows that the highest power of 2 dividing  $T(2^n + 1)$  is  $2^n - 1$ .  $\square$